

# TYLENOL

## DOSING FOR HEALTHCARE PROFESSIONALS

Use this chart to determine the proper dose of **TYLENOL**® for your patient. If possible, use weight to dose; otherwise, use age.<sup>1</sup>

<sup>1</sup>For pediatric products only.

Recommended dose of **TYLENOL**® from your healthcare professional:

\_\_\_\_\_ dose of \_\_\_\_\_ product every \_\_\_\_\_ hours for \_\_\_\_\_ days

		Dosage								
Visit <a href="http://Tylenol.com">Tylenol.com</a> for additional product information		Infants' <b>TYLENOL</b> ® Oral Suspension*	Children's <b>TYLENOL</b> ® Oral Suspension*	Children's <b>TYLENOL</b> ® Chewables*	Children's <b>TYLENOL</b> ® Dissolve Packs	<b>TYLENOL</b> ® Regular Strength Tablets & Liquid Gels	<b>TYLENOL</b> ® Extra Strength Caplets, Coated Tablets, & Rapid Release Gels, & Dissolve Packs	<b>TYLENOL</b> ® For Children + Adults Family*		
Weight (lbs)	Age	Active ingredient: acetaminophen 160 mg (in each 5 mL)	Active ingredient: acetaminophen 160 mg (in each 5 mL)	Active ingredient: acetaminophen 160 mg (in each chewable tablet)	Active ingredient: acetaminophen 160 mg (in each powder)	Active ingredient: acetaminophen 325 mg (in each tablet/liquid gel)	Active ingredient: acetaminophen 500 mg (in each caplet/tablet/powder/gelcap)	Active ingredient: acetaminophen 160 mg (in each 5 mL)		
6-11 lbs	0-3 mos	1.25 mL	The markings on the included dosing cup correspond to the doses listed below. This cup is designed for dosing children ages 2 - 11 years.		<b>Do not use</b>			The markings on the included dosing cup correspond to the doses listed below. This cup is designed for dosing ages 2 years and older.	5 mL	
12-17 lbs	4-11 mos	2.5 mL							7.5 mL	
18-23 lbs	12-23 mos	3.75 mL							10 mL	
24-35 lbs	2-3 yrs	5 mL	5 mL	1 tablet					10 mL	Repeat dose every 4 hours while symptoms last.
36-47 lbs	4-5 yrs		7.5 mL	1 1/2 tablets					12.5 mL	Do NOT give more than 5 doses in 24 hours.
48-59 lbs	6-8 yrs	The markings on the included syringe correspond to the doses listed above. This syringe is designed for dosing children ages 0 - 3 yrs.	10 mL	2 tablets	2 powders				20 mL	
60-71 lbs	9-10 yrs		12.5 mL	2 1/2 tablets	2 powders					
72-95 lbs	11 yrs		15 mL	3 tablets	3 powders					
	<b>Adults and children 12 years and over</b>									

\* The OTC label says "ask a doctor" for dosing of children under 2 years. All Infants' **TYLENOL**® and Children's **TYLENOL**® products have the same strength of acetaminophen: 160 mg (in each 5 mL, tablet, or powder).

**Remind parents & caregivers:**

Read & follow the label on all **TYLENOL**® products. Repeat dose every 4 hours while symptoms last. Do NOT give more than 5 doses in 24 hours. Do NOT use with any other product containing acetaminophen. Use only the dosing device (syringe or dosing cup) that came with the product unless directed by a healthcare provider.

**Remind patients:**

Read & follow the label on all **TYLENOL**® products. Do NOT use with any other product containing acetaminophen. Wait the right amount of time between doses

# Motrin

## DOSING FOR HEALTHCARE PROFESSIONALS

Use this chart to determine the proper dose of MOTRIN<sup>®</sup> for your patient. If possible, use weight to dose; otherwise, use age.<sup>1</sup>

<sup>1</sup>For pediatric products only.

Visit [Motrin.com](http://Motrin.com) for additional product information

mL = milliliter

— dose of — product every — hours for — days

		Dosage					MOTRIN <sup>®</sup> IB Caplets			MOTRIN <sup>®</sup> IB Liquid Gels			MOTRIN <sup>®</sup> IB Migraine Liquid Gels		
Weight (lbs)	Age	Infants <sup>1</sup> MOTRIN <sup>®</sup> Concentrated Drops	Children's MOTRIN <sup>®</sup> Oral Suspension	Children's MOTRIN <sup>®</sup> Chewables	MOTRIN <sup>®</sup> IB Caplets	MOTRIN <sup>®</sup> IB Liquid Gels	MOTRIN <sup>®</sup> IB Liquid Gels	MOTRIN <sup>®</sup> IB Migraine Liquid Gels	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) *Nonsteroidal anti-inflammatory drug.
		Active ingredient: ibuprofen (NSAID)* 50 mg (in each 1.25 mL) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 100 mg (in each 5 mL) *Nonsteroidal anti-inflammatory drug.	Chew or crush tablets completely before swallowing Active ingredient: ibuprofen (NSAID)* 100 mg (in each chewable tablet) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 200 mg (in each caplet) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) *Nonsteroidal anti-inflammatory drug.	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) *Nonsteroidal anti-inflammatory drug.							
12-17 lbs	6-11 mos	1.25 mL			1 tablet										
18-23 lbs	12-23 mos	1.875 mL			1½ tablets										
24-35 lbs	2-3 years				2 tablets										
36-47 lbs	4-5 years				2½ tablets										
48-59 lbs	6-8 yrs				3 tablets										
60-71 lbs	9-10 yrs														
72-95 lbs	11 yrs														

**Remind parents & caregivers:**  
Read & follow the label on all MOTRIN<sup>®</sup> products.  
If needed, repeat dose every 6-8 hours while symptoms last.  
Do NOT give more than 4 doses in 24 hours.  
Use only the dosing device (syringe or dosing cup) that came with the product.  
\*Nonsteroidal anti-inflammatory drug  
Use product only as directed.

**Important Instructions For Proper Use:**  
Always read and follow the label on all MOTRIN<sup>®</sup> products. Keep carton.  
Wait the right amount of time between doses  
If you have any questions, ask your healthcare professional which product may be right for you  
\*Nonsteroidal anti-inflammatory drug

**OTC LABEL STATES:**  
Adults:  
Take 2 capsules with a glass of water; if symptoms persist or worsen, ask your healthcare professional  
**Under 18 years of age:**  
Ask a doctor  
Do NOT take more than 2 capsules in 24 hours unless directed by a doctor

**OTC LABEL STATES:**  
Adults and children 12 years and older:  
Take 1 capsule every 4 to 6 hours while symptoms persist; if pain or fever does not respond to 1 capsule, 2 capsules may be used  
Children under 12 years:  
Ask a doctor  
Do NOT take more than 6 capsules in 24 hours unless directed by a doctor

**OTC LABEL STATES:**  
Adults and children 12 years and older:  
Take 1 caplet every 4 to 6 hours while symptoms persist; if pain or fever does not respond to 1 caplet, 2 caplets may be used  
Children under 12 years:  
Ask a doctor  
Do NOT take more than 6 caplets in 24 hours unless directed by a doctor

Recommended dose of MOTRIN<sup>®</sup> from your healthcare professional: